

Table 8. Percent distribution of nonfatal occupational injuries and illnesses involving days away from work¹ by selected worker characteristics and number of days away from work, Private industry, South Carolina, 2011

| Characteristic | Percent of cases involving | | | | | | | | Median days away from work |
|---|----------------------------|-------|--------|-------------|--------------|---------------|---------------|-----------------|----------------------------|
| | Total cases | 1 day | 2 days | 3 to 5 days | 6 to 10 days | 11 to 20 days | 21 to 30 days | 31 days or more | |
| Total | 100.0 | 18.5 | 10.0 | 18.3 | 12.6 | 7.9 | 3.9 | 28.7 | 6 |
| Gender: | | | | | | | | | |
| Male | 100.0 | 19.6 | 8.8 | 16.9 | 11.2 | 8.3 | 4.6 | 30.7 | 7 |
| Female | 100.0 | 16.5 | 12.8 | 21.0 | 15.7 | 7.2 | 2.4 | 24.5 | 5 |
| Age: | | | | | | | | | |
| 14 - 15 | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| 16 - 19 | 100.0 | 15.4 | 7.7 | 15.4 | 42.3 | -- | -- | 11.5 | 10 |
| 20 - 24 | 100.0 | 24.7 | 17.1 | 17.1 | 15.2 | 8.2 | 1.3 | 16.5 | 3 |
| 25 - 34 | 100.0 | 20.3 | 8.3 | 27.1 | 8.6 | 7.5 | 2.6 | 25.2 | 5 |
| 35 - 44 | 100.0 | 17.0 | 10.3 | 14.3 | 14.3 | 6.7 | 6.7 | 30.5 | 8 |
| 45 - 54 | 100.0 | 14.0 | 10.1 | 14.4 | 8.9 | 9.7 | 3.9 | 38.9 | 14 |
| 55 - 64 | 100.0 | 22.4 | 8.0 | 9.2 | 17.2 | 9.2 | 2.9 | 30.5 | 8 |
| 65 and over | 100.0 | 12.3 | -- | 43.9 | 5.3 | -- | 8.8 | 26.3 | 5 |
| Length of service with employer: | | | | | | | | | |
| Less than 3 months | 100.0 | 19.4 | 15.4 | 26.9 | 9.1 | 9.1 | 1.7 | 18.3 | 4 |
| 3 - 11 months | 100.0 | 16.7 | 9.7 | 20.2 | 12.1 | 8.2 | 4.7 | 28.0 | 6 |
| 1 - 5 years | 100.0 | 20.7 | 8.5 | 16.3 | 15.7 | 4.8 | 3.1 | 31.2 | 6 |
| 5 years or more | 100.0 | 16.8 | 9.7 | 14.2 | 9.7 | 12.3 | 5.6 | 31.7 | 10 |
| Race or ethnic origin ² : | | | | | | | | | |
| White | 100.0 | 22.2 | 7.6 | 22.0 | 12.9 | 6.9 | 3.9 | 24.7 | 5 |
| Hispanic or Latino | 100.0 | 9.5 | 15.9 | 7.9 | 4.8 | 3.2 | -- | 58.7 | 35 |
| Black or African American | 100.0 | 15.1 | 14.7 | 14.2 | 16.4 | 7.8 | 4.3 | 28.0 | 6 |
| Asian | 100.0 | -- | -- | -- | -- | -- | -- | -- | 7 |
| American Indian or Alaska Native | 100.0 | -- | -- | -- | -- | -- | -- | -- | 59 |
| Native Hawaiian or Other Pacific Islander | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| Multi-race | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| Hispanic and other | -- | -- | -- | -- | -- | -- | -- | -- | -- |

Table 8. Percent distribution of nonfatal occupational injuries and illnesses involving days away from work¹ by selected worker characteristics and number of days away from work, Private industry, South Carolina, 2011

| Characteristic | Percent of cases involving | | | | | | | | Median days away from work |
|----------------|----------------------------|-------|--------|-------------|--------------|---------------|---------------|-----------------|----------------------------|
| | Total cases | 1 day | 2 days | 3 to 5 days | 6 to 10 days | 11 to 20 days | 21 to 30 days | 31 days or more | |

¹ Days-away-from-work cases include those that result in days away from work with or without job transfer or restriction.

² Race and ethnicity data do not add to total. Some cases may be counted as both Multi-race and "Hispanic and other" race.

NOTE: Dashes indicate data that are not available. Because of rounding and data exclusion of nonclassifiable responses, data may not sum to the totals.

SOURCE: Bureau of Labor Statistics, U. S. Department of Labor, Survey of Occupational Injuries and Illnesses in cooperation with participating state agencies, November 15, 2013