

Table 8. Percent distribution of nonfatal occupational injuries and illnesses involving days away from work¹ by selected worker characteristics and number of days away from work, Private industry, South Carolina, 2013

| Characteristic | Percent of cases involving | | | | | | | | Median days away from work |
|---|----------------------------|-------|--------|-------------|--------------|---------------|---------------|-----------------|----------------------------|
| | Total cases | 1 day | 2 days | 3 to 5 days | 6 to 10 days | 11 to 20 days | 21 to 30 days | 31 days or more | |
| Total | 100.0 | 12.4 | 9.2 | 15.3 | 10.9 | 10.3 | 6.3 | 35.5 | 13 |
| Gender: | | | | | | | | | |
| Male | 100.0 | 10.9 | 7.9 | 14.1 | 10.5 | 11.1 | 6.1 | 39.4 | 15 |
| Female | 100.0 | 15.5 | 12.2 | 17.5 | 12.0 | 8.7 | 6.7 | 27.4 | 7 |
| Age: | | | | | | | | | |
| 14 - 15 | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| 16 - 19 | 100.0 | 26.7 | 6.7 | 33.3 | 6.7 | -- | -- | 23.3 | 5 |
| 20 - 24 | 100.0 | 12.1 | 13.2 | 15.4 | 11.0 | 12.1 | 9.9 | 26.4 | 10 |
| 25 - 34 | 100.0 | 21.3 | 12.4 | 15.7 | 16.9 | 12.4 | 7.9 | 13.5 | 6 |
| 35 - 44 | 100.0 | 10.0 | 12.6 | 12.6 | 13.4 | 6.9 | 7.8 | 36.8 | 11 |
| 45 - 54 | 100.0 | 10.0 | 5.7 | 13.0 | 7.3 | 12.7 | 4.8 | 46.2 | 22 |
| 55 - 64 | 100.0 | 9.7 | 6.2 | 17.2 | 9.0 | 9.7 | 4.8 | 42.8 | 19 |
| 65 and over | 100.0 | -- | 9.1 | 31.8 | 9.1 | -- | -- | 40.9 | 10 |
| Length of service with employer: | | | | | | | | | |
| Less than 3 months | 100.0 | 7.6 | 6.9 | 13.9 | 12.5 | 6.3 | 3.5 | 50.0 | 30 |
| 3 - 11 months | 100.0 | 13.9 | 10.9 | 11.7 | 12.2 | 6.5 | 8.3 | 36.5 | 11 |
| 1 - 5 years | 100.0 | 14.2 | 7.2 | 15.7 | 9.6 | 11.4 | 6.6 | 35.5 | 14 |
| 5 years or more | 100.0 | 12.0 | 11.4 | 17.9 | 10.8 | 13.9 | 6.2 | 28.1 | 9 |
| Race or ethnic origin ² : | | | | | | | | | |
| White | 100.0 | 13.6 | 12.6 | 15.2 | 11.0 | 10.2 | 8.1 | 29.1 | 8 |
| Hispanic or Latino | 100.0 | 2.9 | 2.9 | 44.9 | 15.9 | 2.9 | 7.2 | 23.2 | 5 |
| Black or African American | 100.0 | 14.3 | 8.5 | 13.3 | 10.9 | 8.5 | 6.1 | 38.4 | 15 |
| Asian | 100.0 | -- | -- | -- | -- | -- | -- | -- | 6 |
| American Indian or Alaska Native | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| Native Hawaiian or Other Pacific Islander | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| Multi-race | 100.0 | -- | -- | -- | -- | -- | -- | -- | 4 |
| Hispanic and other | -- | -- | -- | -- | -- | -- | -- | -- | -- |

Table 8. Percent distribution of nonfatal occupational injuries and illnesses involving days away from work¹ by selected worker characteristics and number of days away from work, Private industry, South Carolina, 2013

| Characteristic | Percent of cases involving | | | | | | | | Median days away from work |
|----------------|----------------------------|-------|--------|-------------|--------------|---------------|---------------|-----------------|----------------------------|
| | Total cases | 1 day | 2 days | 3 to 5 days | 6 to 10 days | 11 to 20 days | 21 to 30 days | 31 days or more | |

¹ Days-away-from-work cases include those that result in days away from work with or without job transfer or restriction.

² Race and ethnicity data do not add to total. Some cases may be counted as both Multi-race and "Hispanic and other" race.

NOTE: Dashes indicate data that are not available. Because of rounding and data exclusion of nonclassifiable responses, data may not sum to the totals.

SOURCE: U.S. Bureau of Labor Statistics, U.S. Department of Labor, Survey of Occupational Injuries and Illnesses in cooperation with participating state agencies, December 05, 2014